



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MICHIGAN

BRAVO WEATHER STIK[®]

EPA Reg. No. 50534-188-100

EPA SLN No. MI-040001

For Control of Diseases on Potato

KEEP OUT OF REACH OF CHILDREN

WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

This label must be in the possession of the user at the time of pesticide application.

RECOMMENDATIONS FOR CONTROL OF DISEASES ON POTATO

BRAVO WEATHER STIK[®] is an excellent disease control agent when used according to label directions for control of a broad spectrum of plant diseases. **BRAVO WEATHER STIK[®]** is recommended for use in programs which are compatible with the principles of Integrated Pest Management (IPM), which include the use of disease resistant crop varieties, cultural practices, pest scouting and disease forecasting systems which reduce unnecessary applications of pesticides.

BRAVO WEATHER STIK[®] is effective for strategic use in programs that attempt to minimize disease resistance to fungicides. Some other fungicides, which are at risk from disease resistance, exhibit a single-site mode of fungicidal action. **BRAVO WEATHER STIK[®]**, with a multi-site mode of action, may be used to delay or prevent the development of resistance to single-site fungicides. Consult with your federal or state Cooperative Extension Service representatives for guidance on the proper use of **BRAVO WEATHER STIK[®]** in programs that seek to minimize the occurrence of disease resistance to other fungicides.

CROP	DISEASE	CASUAL AGENT	RATE PER ACRE (lbs. a.i./A)
Potato	Late Blight Early Blight Botrytis vine rot	Phytophthora infestans Alternaria solani Botrytis cinerea	Low rate: 3/4 pt (0.6) then, Higher rates: 1 to 1 ½ pt (0.75 to 1.125)

APPLICATION DIRECTIONS

Begin applications of **BRAVO WEATHER STIK** at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals.

Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occurs:

- Vines close between rows;
- Late blight forecasting measures 18 disease severity values (DSV); or
- The crop reaches 300 P-days.

Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe. A maximum of 21-1/2 pints of **BRAVO WEATHER STIK** (16 pounds a.i.) per acre may be used on long-season varieties of potato during each growing season. **DO NOT** apply within 7 days of harvest.

BRAVO WEATHER STIK® may be applied through sprinkler irrigation equipment (solid set, portable wheel move, motorized lateral move or center pivot systems only). **DO NOT** exceed a 10 day interval between applications when using this technique. See calibration directions on **BRAVO WEATHER STIK®** label.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash waters. **DO NOT** apply when weather conditions favor drift from treated areas. **APPLY ONLY TO AREAS SPECIFIED ON LABEL.**

BRAVO® and WEATHER STIK® are registered trademarks of GB BIOSCIENCES CORPORATION.

24(c) Sub-Registrant:
Syngenta Crop Protection, Inc.
Greensboro, NC 27419

EPA SLN No. MI-040001
Expiration Date: January 10, 2009

MI0188018AA0104

MIBravoWeatherStikPotatoSLNLbl-SYN 1-28-05